



Nutritional Composition of *Mucuna pruriens* var. *utilis* (Velvet bean) Seeds Grown in Nanyumbu District, Tanzania

Florence Boniface^{1*}, Stephen I. Nnungu² and Washa B. Washa¹

¹Mkwawa University College of Education, University of Dar es Salaam, P. O. Box 2513 Iringa, Tanzania

²Department of Botany, College of Natural and Applied Sciences, University of Dar es Salaam, P. O. Box 35060 Dar es Salaam, Tanzania

Emails: florencenyungu3@gmail.com

*Corresponding author

Received 8th Aug. 2024, Reviewed 30th Aug., Accepted 15th Oct., Published 30th Nov. 2024

<https://dx.doi.org/10.4314/tjs.v50i4.1>

Abstract

In tropical countries, *Mucuna* seeds have been utilized to promote better health and ensure a stable food supply. The use of *Mucuna pruriens* var. *utilis* seeds in Tanzania has prompted an evaluation of the nutritional composition of two common cultivars grown in the Nanyumbu district. *Mucuna* seed samples (black-reddish brown and black-greyish) were collected and analyzed for their nutritional content using standard extraction and spectrophotometric techniques. Data analysis was conducted using GraphPad InStat 3.0 software, with a t-test employed to compare means at a 95% confidence level. The results indicated that the differences in nutritional parameters between the black-reddish brown and black-greyish *Mucuna* seeds were not significant ($P > 0.05$), except for crude lipid and carbohydrate content. The protein, lipid, carbohydrate, fiber, ash, vitamin C, beta carotene, lycopene, and energy content ranged from 48.13% to 46.50%, 2.13% to 1.39%, 26.75% to 23.12%, 15.99% to 14.51%, 3.23% to 3.21%, 0.98 to 0.84 mg/g, 0.011 to 0.004 mg/100 mg, 0.006 to 0.0058 mg/100 mg, and 1303.7 to 1243.1 kJ/100 g, respectively. Both cultivars of *Mucuna* seeds exhibit dietary qualities that can be explored as alternative food sources for the malnourished population in the country.

Keywords: Black-greyish seeds, Black-reddish brown seeds, Cultivars, Nutritional quality

Introduction

Velvet bean, *M. pruriens* L., commonly known as ‘Upupu,’ is an annual climbing legume from the Fabaceae family (Buckles 1995). This plant species originated in Eastern India and China and has since spread throughout tropical regions worldwide (Wilmot-Dear 1984, Suryawanshi et al. 2020). *Mucuna* is widely recognized for its nutritional and bioactive contributions to human health (Timsina 2018, Pathania et al. 2020). During periods of food scarcity, *Mucuna* seeds serve as a crucial safeguard against famine (Huisden 2008). Various ethnic groups and undernourished populations have traditionally used these

seeds as a food source (Janardhanan et al. 2003; Lampariello et al. 2012). Studies have shown that *Mucuna* seeds can help alleviate malnutrition in infants when consumed by nursing mothers (Huisden 2008). Currently, the species is cultivated in Western Tanzania as green manure (Matata et al. 2017). In southern Tanzania, however, *Mucuna* is often undervalued and primarily used as food (Constantine et al. 2020, Sakamoto et al. 2020). Given that the plant’s chemical composition can vary with growing conditions, nutritional analyses of *Mucuna* seeds from Nanyumbu, Tanzania, are warranted (Kalidas and Mohan 2011).

The appeal of *Mucuna* as a functional food lies in its high nutritional and potential phytochemical content (Encalada and Campos 2021). *Mucuna* seeds demonstrate long viability, pest resistance, and high germination rates even in challenging conditions (Siddhuraju et al. 2000). *Mucuna* has a high yielding capacity across various environmental conditions (Buckles 1995; Carew and Gernat 2016), thriving best in warm, moist climates with sandy or loamy soils at altitudes of about 1600 m (Buckles 1995). The plant can tolerate drought, low soil fertility, and high acidity (Correia et al. 2014). The life cycle of common varieties typically ranges from 160 to 170 days, with flowering occurring between the 59th and 66th days (Sathyanarayana et al. 2012).

As the plant adapts to environmental stresses, it accumulates antinutritional substances that may impact its nutritional value and could lead to adverse effects when ingested (Lorenzetti et al. 2010, Banti & Bajo 2020). Moreover, the presence of high levels of these factors limits the acceptance of *Mucuna*. The pods contain brown hairs with mucunain protein, which can cause severe itching upon contact with skin (Mosissa et al. 2021). Phytochemicals identified as potential antinutritional factors in *Mucuna* seeds include L-DOPA, phenolics, flavonoids, and ursolic acid (Kala et al. 2010, Rakesh and Praveen 2020). L-DOPA can be harmful if consumed in improperly processed seeds (Kosower and Kosower 1967, Lorenzetti et al. 2010). However, when seeds are properly processed, they provide beneficial bioactive compounds associated with pharmacological and antioxidant properties (Rai et al. 2017; Rai et al. 2019). This highlights the necessity of including *Mucuna* in the diet, as these compounds cannot be synthesized in the human body (Kumar and Pandey 2013).

The accessibility and high yield of the two common cultivars of *Mucuna* seeds in the Nanyumbu district ensure a reliable food supply throughout the year. Despite its potential, there is limited information on the nutritional composition of *M. pruriens* var. *utilis* seeds cultivated in Nanyumbu, Tanzania. This study aims to evaluate the nutritional composition of two common cultivars of *M. pruriens* var. *utilis* seeds and compare the nutritional parameters between black-reddish brown and black-greyish *Mucuna* seeds grown in the Nanyumbu district.

Materials and Methods

Description of the study area

The study was conducted in Nanyumbu District, covering four villages: Namatumbusi, Mikangaula, Kilosa, and Chang'ombe, which lie within the grid range of 10°49'54"S to 10°50'30"S and 38°38'36"E to 38°41'24"E. The four villages are situated at elevations between 330 and 340 meters, within Nanyumbu district, which lies in the lowlands at altitudes ranging from 0 to 501.343 meters above sea level. Two cultivars of *M. pruriens* var. *utilis*, with black-reddish brown and black-greyish seed coats, exhibit high performance in the study area. Nanyumbu district receives an average annual rainfall of 800 mm, with a single rainy season that lasts from November to May. The mean temperature is 27 °C during hot periods and 24 °C during cool periods. Figure 1 presents a map of Nanyumbu district, indicating the four sampling microsites of the two common cultivars of *M. pruriens* var. *utilis*. The natural soil of Nanyumbu district is fertile and composed of sandy, loamy, gritty, and gravelly structures.




Figure 1: A map of Nanyumbu District indicating four sampling sites

Collection and preparation of seed sample

Two seed samples of *M. pruriens* var. *utilis* (black-reddish brown and black-greyish) were collected from four microsites as mature dry pods during harvesting period of July, 2022 by adopting procedure described by Ade-Omowaye et al. (2015). With the help of keys by Gillett et al. (1971) the seeds were botanically identified. Dry pods were threshed to remove seeds and thorough winnowed. A 500 g of each seed sample were collected from every village, packed in a polythene bags, well labelled and transferred to the Botany department, University of Dar es Salaam for nutritional analysis. Sample preparation was done following the procedure described by Nwaoguikpe et al. (2011).

Laboratory analysis

For every cultivar, collected seeds sample were separately grounded using a laboratory mill into fine powder, labelled and kept in refrigerator at 4 °C for nutritional analysis (Siddiq et al. 2010). Each accession of *Mucuna* seeds was analysed in triplicates for crude protein, crude lipid, crude fiber, total

ash, vitamin C, β -carotene, lycopene and carbohydrate. For chemical extraction this study adopted the procedures described by James (1995). The study employed the use of UV-Visible spectrophotometer (Jenway 6560, Limited, U.K) in all readings. Chemical analysis to determine nutritional composition of each sample of *Mucuna* seeds was done using standard procedure.

Determination of protein content

Crude proteins derived from *Mucuna* seeds was done based on the total nitrogen (N) content through micro-Kjeldahl method as described by Burman et al. (1996). An extract was made from 0.1 g of raw sample were digested in micro-Kjeldahl at 380 to 340 °C and oxidized by 97% concentrated sulphuric acid under copper catalyst and potassium sulphate that raised boiling point of a mixture. The cold digests and standard (NH₄Cl) was mixed with a combined reagent (0.35 g of sodium potassium tartrate, 0.17 g sodium salicylate, 0.005 g nitroprusside and 0.4 ml of 50% sodium hydroxide) and 4 ml of 0.15% hypochlorite solution at the ratio of

5:40:4 ml which then reduced to 1:8:0.8 ml. Both treatments and standard read absorbance at 660 nm. A created calibration curve was used to calculate concentration of total nitrogen [x] from the standard using an equation:

$$[x] = (y - b)/m$$

Where x = concentration of standard ammonium chloride, y = absorbances of sample extract, b = y-intercept and m = slope. A [x] was used to obtain the percentage composition of total nitrogen (%N) by equation;

$$\%N = (\text{concentration (x)} \times 100)/10 \times 8 \times 0.05$$

Where 10, 8, 0.05 were dilution factors used for nitrogen

Then, protein (P) content was obtained by the equation $P = \%N \times 6.25$ (Tomé et al. 2022).

Determination of crude lipid content

Crude lipid content was determined using chloroform-methanol method (Nielsen 1998). An extract was obtained by dissolving 3 g of a grounded sample in chloroform, 78% methanol and distilled water of 1:2:0.8 ratio. A chloroform was added in a mixture to double its volume to the ratio of 2:2:0.8 and the content was homogenized for 30 seconds. Finally, distilled water was added to double its volume making a ratio of 2:2:1.6. The content was then filtered through a whatman No. 1 filter paper. The layered filtrate was separate and an eluted upper layer was placed in oven at 40 °C for 24 hours to evaporate chloroform leaving the lipid extract. Residual weight of lipid content was obtained by subtracting the weight of a beaker containing lipid with that of an empty beaker. Percentage composition of crude lipid was obtained using the equation below

$$\% \text{ Crude lipids} = [(\text{volume of solvent} \times \text{residual weight}) / \text{aliquot} \times 15] \times 100$$

Where 15 is a dilution factor

Determination of carbohydrate content

Carbohydrate was determined by anthrone calorimetric method (Maness 2010). Anthrone reagent (760 ml concentrated sulphuric acid, 330 ml distilled, 1 g of Anthrone and 1 g of Thiourea) treated in cool and dark place for 30 min. A 2 g of grounded seeds was dissolved in hot water and later allowed to cool then filtered. Each of filtrates,

standard (A freshly prepared sugar solution of 0.25 g/l) and water blank in boiling tubes were treated in cool and dark place. A 10 ml Anthrone reagent was added while the tubes immersed in running cold water. The contents in tubes were warmed while placed in dark and boiled, thereafter cooled. Standards, sample extracts and a blank absorbances were read at 625 nm. A prepared calibration graph was used to obtain milligram of glucose in the sample of four aliquots which was expressed as glucose equivalent percent. Carbohydrate concentration was calculated using an equation;

$$Y = m[x] + b$$

Where x = concentration of glucose in mg/ml, Y = absorbances of the sample, b = y-intercept and m = slope or a gradient factor.

The percentage composition of was determined using equation;

$$\% \text{ Carbohydrate} = (x \times 30)/10 \times 2 \times 0.05$$

Where 2 was a mass of sample, 30, 10 and 0.05 were used as dilution factors for carbohydrates. The validity of results was justified with correlation coefficient (R^2).

Determination of energy value

The calorific equation was used to estimate energy value of *Mucuna* seeds by multiplying the percentage of crude protein, crude lipid and crude carbohydrate by 16.7, 37.7 and 16.7, respectively (Siddhuraju et al. 1996).

Determination of crude fibre

Procedure described by James et al. (2020) was adopted to determine crude fibre content in *Mucuna* seeds. A 1 g of sample was acidified by boiling with HCl (1%) then, contents were filtered and boiled with an alkali (NaOH). The sample residue was dried, cooled and weighed. Dry residues were ashed in a muffle furnace at 550 °C then cooled and re-weighed. Percentage of weight loss of an incinerated residue was considered as a weight of crude fiber content as calculated using an equation:

$$\% \text{ Crude fibre} = [(\text{weight of sample} - \text{weight of shed}) / \text{weight of original sample}] \times 100$$

Determination of total ash

Total ash of the *Mucuna* seeds were determined following procedure described by James et al. (2020). Grounded samples of 5 g was incinerated in a muffle furnace at 500 °C.

The percentage weight of ashes was calculated using the equation;

$$\% \text{ Ash} = (\text{weight of ash} / \text{weight of sample}) \times 100$$

Determination of vitamin C

Vitamin C was extracted from 2 g of grounded *Mucuna* seeds and determined by 2, 4-Dinitrophenylhydrazine assay as explained by Desai and Desai (2019). A sample was dissolved in 5 ml of 4% oxalic acid then homogenized and centrifuged at 5000 rpm for 10 min. Supernatant contents were filtered through a whatman no. 1 filter paper. The standard was prepared in a ratio of 1:2 (ascorbic acid-distilled water). Each of sample aliquot and standard were oxidized with bromine water. In every content of sample and standard a 1 ml of 2, 4-Dinitrophenylhydrazine reagent was added followed by two drops of thiourea water and incubated at 37 °C for 3 hours in dark cupboard. Both contents of sample and standard were crystalised which were dissolved in 5 ml of 80% sulphuric acid. The resulted red colour solutions read absorbance at 540 nm. The graph created was used to calculate concentration of ascorbic acid (mg/l) from the equation;

$$C \text{ (mg/l)} = (A_{540} - y) / m$$

Where C = concentration of the standard, A_{540} = Absorbance of the sample read at 540 nm, y = y-intercept, m = gradient factor
Finally ascorbic acid content in mg/g of sample dry weight were calculated by a formula;

$$\text{Concentration (mg/g)} = \text{Concentration of standard (mg/l)} \times 10 \text{ factor}$$

Determination of beta-carotene and lycopene

A petroleum ether method described by Barros *et al.* (2007) was used to determine β -carotene and lycopene content of *Mucuna* seeds. A 2 g of grounded *Mucuna* seeds was dissolved in 5 ml of 0.5 N ethanoic potassium hydroxide solution and 15 ml of petroleum ether then filtered. An upper petroleum ether layer formed by addition of distilled water were eluted and evaporated to dryness on a hot plate at 70 °C. A 100 g of cool dried ether extract was dissolved in 10 ml acetone-hexane mixture (4:6) and filtered through

whatman No.1 filter paper. The absorbance of the filtrates was measured at 663, 505 and 453 nm. The content of beta carotene and lycopene were calculated according to the equations below suggested by Barros *et al.* (2007).

$$\text{Beta carotene (mg/100mg)} = (0.216 \times A_{663}) - (0.304 \times A_{505}) + (0.452 \times A_{453})$$

$$\text{Lycopene (mg/100mg)} = (-0.0458 \times A_{453}) + (0.372 \times A_{505}) - (0.0806 \times A_{663}).$$

Data analysis

Data were analyzed using GraphPad InStat software version 3.0 (Motulsky *et al.* 2003). Descriptive statistics were presented in tables. Variations in means for each tested parameter were compared between the black-reddish brown and black-greyish *Mucuna* seeds using an unpaired t-test at a 95% confidence level. All determinations were carried out in triplicate, and data were expressed as mean \pm standard deviation.

Results and Discussion

Nutritional composition

Research on the nutritional composition of neglected food plants plays a significant role in enhancing the food industry. *M. pruriens* var. *utilis* seeds exhibit nutritional potential that promotes their use as a functional food source for undernourished populations. The nutritional composition of two cultivars of *M. pruriens* var. *utilis* seeds (black-reddish brown and black-greyish) is summarized in Table 1. The findings indicate that the nutritional composition of black-reddish brown and black-greyish seeds did not differ significantly ($P > 0.05$) in all tested parameters, except for crude lipid and carbohydrate content.

Protein content

Both cultivars of *M. pruriens* var. *utilis* grown in the Nanyumbu district recorded high protein levels ranging from 46.50% to 48.13%. These levels are significantly higher than the 43.12% to 43.40% reported for *M. pruriens* seeds (Renata *et al.* 2015). Additionally, the protein content is nearly double that of the white variety (29.53%) and black variety (28.75%) of *M. pruriens* (Daffodil *et al.* 2016). Lower protein levels were also reported for the white variety

(28.82%) and black seeds (26.26%) (Kalidass and Mahapatra 2014). The protein content in *Mucuna* seeds is about twice that found in commonly consumed legume seeds in the tropics, such as *Phaseolus vulgaris* (common bean) at 19.45% (Grela et al. 2017). Even lower protein levels have been noted in bambaranut, red bean, pigeon pea, cowpea, and groundnut (James et al. 2020). In comparison, the protein content of the *Mucuna* seeds in this study exceeds the 41.81% reported for soybean seeds (Mnembuka and Eggum 1995). The protein levels found in *Mucuna* seeds exceed the daily protein needs for adults, which range from 10% to 35% as recommended by the USA National Research Council (1989).

Carbohydrate content

The black-reddish brown seeds in this study exhibited a higher carbohydrate level (26.747%) compared to the black-greyish seeds (23.12%). The carbohydrate content in the black-reddish brown seeds aligns with the black (27.19%) and white (28.27%) varieties of *M. pruriens* var. *utilis* (Siddhuraju and Becker 2001). In contrast, the black-greyish seeds showed a lower carbohydrate content (23.121%) than the 28% found in *Mucuna* bean seeds (Siddhuraju and Becker 2005). These findings are consistent with carbohydrate levels in other pulses, including pigeon pea (23.66%) (Ade-Omowaye et al. 2015). The carbohydrate content is significantly higher than that of groundnut (7.34%) (James et al. 2020), soybean (1.33%), and winged bean (3.00%) (Mnembuka and Eggum 1995). However, both cultivars recorded lower carbohydrate levels compared to other popular legumes, such as chickpea (31.8%) and green gram (39.9%) (Bravo et al. 1999). Other legumes like green gram (38.54%), bambaranut (45.02%), pigeon pea (40.53%), field bean (36.30%), and cowpea (41.34%) (Mnembuka and Eggum 1995) also displayed higher carbohydrate content than the current findings. The carbohydrate levels of 26.747% to 23.121% in both studied *Mucuna* seeds exceed the recommended intake of 21.2% for managing cardiovascular disease risk (Shan et al. 2019).

Crude lipid content

The black-greyish seeds exhibited a higher lipid content (2.13%) compared to the black-reddish brown seeds (1.39%). The lipid content range of 2.13% to 1.39% in all cultivars of *M. pruriens* var. *utilis* aligns with the 2.69% found in *M. pruriens* seeds (Nwaoguikpe et al. 2011), but is lower than that reported for the white and black varieties (Daffodil et al. 2016, Kalidass and Mahapatra 2014, Tresina and Mohan 2013). Similar lipid levels were noted in other preferred legumes, such as *P. vulgaris* (1.190%) (Shimelis and Rakshit 2005), kidney bean (1.4%), lima bean (1.3%), pigeon pea (2.8%), and jack bean (2.5%) (Apata and Ologhobo 1994). Some commonly cultivated legumes, such as bambaranut, red bean, pigeon pea, and cowpea, reported higher crude lipid contents (5.87%, 6.50%, 6.73%, and 6.75%, respectively) (James et al. 2020). The crude lipid content does not categorize *M. pruriens* var. *utilis* grown in the Nanyumbu district as an oil-rich legume, especially when compared to groundnuts (34.63%) (James et al. 2020), soybeans (22.88%), and winged beans (21.75%) (Mnembuka and Eggum 1995).

Energy value

The energy value estimated for both cultivars ranged from 1243.1 to 1303.7 kJ/100 g (dry matter). These values are lower than the 1570.34 to 1607.66 kJ/100 g reported by Renata et al. (2015) for *M. pruriens* seeds, but higher than the 572.95 kJ/100 g of *M. pruriens* var. *utilis* (Rane et al. 2019). Compared to other preferred legumes, Mnembuka and Eggum (1995) reported higher energy values in soybean (2383.21 kJ/100 g), green gram (1945.14 kJ/100 g), bambara nut (1973.17 kJ/100 g), pigeon pea (1881.55 kJ/100 g), field bean (1877.78 kJ/100 g), cowpea (1925.48 kJ/100 g), and winged bean (2372.33 kJ/100 g).

Crude fiber content

The findings indicate a crude fiber content ranging from 14.51% to 15.99%, which is higher than the levels reported for the white and black varieties of *M. pruriens* var. *utilis* (Daffodil et al. 2016, Kalidass and Mahapatra 2014, Tresina and Mohan 2013). These

findings also surpass the fiber content of common pea, common bean, and chickpea (4.571%, 4.434%, and 2.216%, respectively) (Grela et al. 2017). Lower fiber content has been reported in bambaranut, red bean, pigeon pea, cowpea, and groundnut (2.23%, 2.50%, 3.13%, 3.16%, and 6.11%, respectively) (James et al. 2020). The fiber content in both cultivars indicates that only 2 g of *Mucuna* pulses would meet the recommended daily fiber intake of 25–30 g for adults (Reynolds et al. 2019). Although fiber does not contribute directly to the nutritive value of foods, it plays a vital role in managing peristalsis and metabolism in the gastrointestinal tract (Daffodil et al. 2016, Tresina and Mohan 2013).

Total ash content

The total ash content of all studied *Mucuna* seeds ranged from 3.21% to 3.23%, aligning with the 3.10% found in *M. pruriens* seeds (Renata et al. 2015) and 3.3% in *Mucuna* accessions (Vadivel and Janardhanan 2000). These values are significantly lower than the 6.47% reported by Alaye et al. (2020) and the 4.94% and 5.30% for the black and white varieties, respectively (Tresina and Mohan, 2013). Comparatively, similar results were observed in chickpea (3.13%) (Grela et al., 2017), jack bean (3.0%) (Apata and Ologhobo 1994), and *P. vulgaris* (3.024%) (Shimelis and Rakshit 2005). Lower levels were reported in common pea (2.746%) (Grela et al., 2017) and lentils (2.8%) (Iqbal et al. 2006). The available ash content

suggests that *Mucuna* seeds contain nutritionally significant mineral components (Kalidas and Mohan 2011; Tresina and Mohan 2013).

Vitamin C content

The vitamin C levels observed in both cultivars ranged from 0.982 to 0.8406 mg/g, significantly higher than the 0.55 and 0.5832 mg/g found in the black and white varieties of *M. pruriens* var. *utilis* (Tresina and Mohan 2013). Lower vitamin C levels were reported in mung gram, black gram, pigeon pea, and cowpea (Tomar et al. 2018). Despite the relatively low levels found in this study, the vitamin C content underscores the potential of *Mucuna* seeds as an important functional food source rich in antioxidants.

Beta-Carotene and Lycopene levels

The mean beta-carotene levels in both cultivars ranged from 0.011 to 0.004 mg/100 mg, higher than the 0.00015 to 0.00346 mg/100 mg reported for *M. flagellipes* (Nwajagu et al. 2021). Lower beta-carotene levels were noted pigeon pea and cowpea (Tomar et al. 2018), as well as in lupin seeds (11.98–50.43 µg/g) (Wang et al. 2008) and soybean seeds (Gebregziabher et al. 2021, 2022). Lycopene levels in this study ranged from 0.0062 to 0.0058 mg/100 mg, similar to the 12.7 mg/100 g (0.0127 mg/100 mg) found in lima beans (Farinde et al. 2017). In this context, *Mucuna* seeds appear to be a promising crop with antioxidant properties capable of combating free radicals.

Table 1: Nutritional composition of two accessions of *Mucuna pruriens* var. *utilis*; black reddish brown and black greyish seeds

Component	<i>Mucuna pruriens</i> var. <i>utilis</i> (Black reddish brown seed coat)	<i>Mucuna pruriens</i> var. <i>utilis</i> (Black greyish seed coat)	t-value	P-value
Crude Protein (%)	48.13±6.89 ^a	46.50±4.89 ^a	0.667	0.5117
Crude Lipid (%)	1.39±0.39 ^b	2.133±0.09 ^a	5.144	0.0001
Carbohydrate (%)	26.75±3.16 ^a	23.12±2.23 ^b	3.247	0.0037
Total Fibre (%)	14.51±1.25 ^a	15.99±2.28 ^a	1.61	0.1297
Total Ash (%)	3.23±0.03 ^a	3.21±0.102 ^a	0.428	0.6752
Vitamin C (mg/g)	0.84±0.26 ^a	0.98±0.007 ^a	1.916	0.0684
β-carotene (mg/100 mg)	0.011±0.02 ^a	0.004±0.003 ^a	1.549	0.1356
Lycopene (mg/100 mg)	0.006±0.005 ^a	0.006±0.003 ^a	0.2704	0.7894
Energy (kJ/100 g DM)	1302.736 ^a	1243.118 ^b		

All values are expressed as a Mean ± SD, n = 3. Mean values in the following row sharing a common letter in superscript are not statistically significant at 95% confidence

level. DM = Digestible and Metabolized energy

Conclusion

The primary constituents of *Mucuna* seeds identified in this study were protein, carbohydrates, and fiber. The black-reddish brown seeds exhibited the highest levels of crude protein, carbohydrates, ash, beta carotene, lycopene, and energy. In contrast, the black-greyish seeds contained higher amounts of crude lipids, crude fiber, and vitamin C. The findings suggest that the nutritional profile of *M. pruriens* var. *utilis* (both black-reddish brown and black-greyish seeds) is comparable to, and even exceeds, that of other *Mucuna* seeds from similar or different varieties. The high protein content in these seeds may help combat protein-energy malnutrition, while the levels of carbohydrates, lipids, and fiber can effectively address cholesterol and cardiovascular issues. Although *Mucuna* seeds contain anti-nutritional factors, these are not expected to pose significant risks to humans and may even provide antioxidant benefits at levels deemed healthy. There is a pressing need for further research on *Mucuna* species, as they possess the potential to serve as functional foods and natural medicines for large populations.

Acknowledgement

We acknowledge the University of Dar es Salaam for material support in completion of this study.

References

- Ade-Omowaye BIO, Tucker G and Smetanska I 2015 Nutritional Potential of nine Underexploited Legumes in Southwest Nigeria. *Int. Food Res. J.*, 22(2): 798–806.
- Alaye S, Layade K, Omole E, Onihunwa J, Joshua D and Akande O 2020 Effects of Different Processing Methods on Proximate Composition of *Mucuna Pruriens*. *Int. J. of Progress. Sci. Technol.* 20(2): 229–233.
- Apata DF and Ologhobo AD 1994 Biochemical Evaluation of Some Nigerian Legume Seeds. *Food Chem.* 49(4): 333–338.
- Banti M and Bajo W 2020 Review on Nutritional Importance and Anti-nutritional Factors of Legumes. *Int. J. Nutr. Food Sci.* 9(6): 138.
- Barros L, Ferreira MJ, Queirós, B Ferreira IR and Baptista P 2007 Total phenols, ascorbic acid, β -carotene and lycopene in Portuguese wild edible mushrooms and their antioxidant activities. *Food Chem.*, 103(2): 413–419.
- Bravo L, Siddhuraju P and Saura-Calixto F. 1999 Composition of underexploited Indian pulses. Comparison with common legumes. *Food chem.* 64(2): 185–192.
- Buckles D 1995 Velvetbean: A “new” plant with a history. *Econ. Bot.* 49(1), 13–25.
- Buurman P, Lagen B Van and Velthorst E J 1996 WfM Manual for soil and water analysis.
- Carew LB and Gernat AG 2016 Velvet Bean (*Mucuna pruriens*) *Feedipedia*.
- Constantine J, Sibuga KP, Shitindi MJ, and Hilberk A 2020 Awareness and Application of Existing Agroecological Practices by Small Holder Farmers in Mvomero and Masasi Districts-Tanzania. *J. Agric. Sci.* 13(1): 30.
- Correia MV, Pereira LR, De Almeida L, Williams RL, Freach J, Nesbitt H and Erskine W 2014 Maize-mucuna (*Mucuna pruriens* (L.) DC) relay intercropping in the lowland tropics of Timor-Leste. *Field Crops Res.* 156: 272–280.
- Daffodil ED, Tresina PS, and Mohan VR 2016 Nutritional and Antinutritional Assessment of *Mucuna pruriens* (L.) DC var. *utilis* (Wall ex. Wight) Bak. Ex Burck and *Mucuna deeringiana* (Bort) Merrill: An Underutilized Tribal Pulse. *Int. Food Res. J.* 23(4): 1501–1513.
- Desai AP and Desai S 2019 UV Spectroscopic Method for Determination of Vitamin C (Ascorbic Acid) Content in Different Fruits in South Gujarat Region. *Int. J. Envir. Sci. Nat. Resour.* 22(2): 41–44.
- Encalada and Campos 2021 *Mucuna pruriens* Fiber: Nutritional, Functional and Biological Properties. *Food Sci. Technol. (Brazil)*, 41(1): 120–126.
- Farinde EO, Obatolu VA and Fasoyiro SB

- 2017 Microbial, Nutritional and Sensory Qualities of Baked Cooked and Steamed Cooked Lima Beans. *Am. J. Food Sci. Technol.*, 5(4), 156–161.
- Gebregziabher, Berhane S, Zhang S, Azam M, Qi J, Agyenim-Boateng KG, Feng Y, Liu Y, Li J, Li B and Sun J 2022 Natural Variation and Geographical Distribution of Feed Carotenoids and Chlorophylls in 1167 Chinese Soybean Accessions. *J. Int. Agric.* 1–21.
- Gebregziabher, Berhane Sibhatu, Zhang, S, Qi J, Azam M, Ghosh S, Feng Y, Huai, Y, Li J, Li B and Sun J 2021 Simultaneous Determination of Carotenoids and Chlorophylls by the HPLC-Uv-Vis Method in Soybean Seeds. *Agronomy* 11(4): 1–15.
- Gillett JB, Polhill R and Verdcourt B 1971 Flora of tropical East Africa. Leguminosae (Part 3) Subfamily Papilionoideae (1); Leguminosae (Part 4) Subfamily Papilionoideae (2). *Flora of Tropical East Africa. Leguminosae (Part 3) Subfamily Papilionoideae (1); Leguminosae (Part 4) Subfamily Papilionoideae (2)*.
- Grela ER, Kiczorowska B, Samolińska W, Matras J, Kiczorowski P, Rybiński W, and Hanczakowska E 2017 Chemical Composition of Leguminous Seeds: Part I—Content of Basic Nutrients, Amino Acids, Phytochemical Compounds, and Antioxidant Activity. *Eur. Food Res. Technol.* 243(8): 1385–1395.
- Huisden CM 2008 Detoxification, Nutritive Value, and Anthelmintic Properties of *Mucuna pruriens*. *Anim. Sci.*, 1–131.
- Iqbal A, Khalil IA, Ateeq N, and Sayyar KM 2006 Nutritional Quality of Important Food Legumes. *Food Chem.*, 97(2), 331–335.
- James CS 1995 Analytical Chemistry of Foods. *Chapman and Hall* <https://doi.org/10.1007/978-1-4615-2165-5>
- James S, Nwabueze TU, Onwuka GI, Ndife J, Usman MA, and A 2020 Chemical and Nutritional Composition of Some Selected Lesser Known Legumes Indigenous to Nigeria. *Heliyon* 6(11): e05497.
- Janardhanan K, Vadivel V and Pugalenth M 2003 Biodiversity in Indian Underexploited/Tribal Pulses. *Kluwer Acad.* 046: 353–405.
- Kala BK, Kalidass C, and Mohan VR 2010 Nutritional and Antinutritional Potential of Five Accessions of a South Indian Tribal Pulse *Mucuna atropurpurea* DC. *Trop. Subtrop. Agroecosystems* 12: 339–352.
- Kalidas and Mohan 2011 Nutritional and Antinutritional Compositional of Itching Bean (*Mucuna pruriens* (L.) DC var. *pruriens*): an Underutilized Tribal Pulse in Western Ghats, Tamil Nadu. *Trop. Subtrop. Agroecosystems* 14: 279–293.
- Kalidass C and Mahapatra AK 2014 Evaluation of the Proximate and Phytochemical Compositions of an Underexploited Legume *Mucuna pruriens* var. *utilis* (Wall ex Wight) L.H. Bailey. *Int. Food Res. J.* 21(1): 303–308.
- Kosower NS and Kosower EM 1967 Does 3, 4-dihydroxyphenylalanine play a part in favism? *Nature* 215(5098): 285–286.
- Kumar S and Pandey AK 2013 Chemistry and Biological Activities of Flavonoids. *Sci. World J.*, 2013: 162750
- Lampariello L, Cortelazzo A, Guerranti R, Sticozzi C and Valacchi G 2012 The Magic Velvet Bean of *Mucuna pruriens*. *J. Tradit. Complem. Med.* 2(4): 331–339.
- Lorenzetti F, Macisaac S, Arnason JT and Buckles D 1998 The Phytochemistry, Toxicology, and Food Potential of Velvet bean (*Mucuna adans.* spp. Fabaceae). In: Cover crops in West Africa (Conference proceedings) 1-3 October 1996, Cotonou, Benin. 1–11.
- Maness N 2010 Extraction and Analysis of Soluble Carbohydrates. *Plant Stress Tolerance* 639: 341–370.
- Matata PZ, Passos AMA, Masolwa LW, Marcolan AL and Ribeiro R da S 2017 Incorporation of Leguminous Cover Crops in Smallholder Cassava-Based Production System in Western Tanzania. *Am. J. Plant Sci.* 08(13): 3490–3501.
- Mnembuka BV and Eggum BO 1995 Comparative Nutritive Value of Winged

- Bean (*Psophocarpus tetragonolobus* (L) DC) and other Legumes Grown in Tanzania. *Plant Foods Hum. Nutr.* 47(4), 333–339.
- Mosissa D, Awas T, Atinafu H and Olana T 2021 Agro-Botanical Characterization of Different Accessions of *Mucuna Melanocarpa* Hochst A Rich. *South Asian Res. J. Biol. Appl. Biosci.* 3(4): 46–50.
- Motulsky HJ, Paige S, Pilkington J and Platt M 2003 InStat3, Guide to Choosing and Interpreting Statistical Tests. 1–126.
- Nielsen SS 1998 Moisture and Total Solids Analysis. In *Instructor's Manual for Food Analysis: Second Edition*.
- Nwajagu IU, Nzelihe HC, Chukwuekezie N E, Abah CR, Umar AT and Anarado CS 2021 Effect of Processing on the Nutrient, Anti-nutrient and Functional Properties of *Mucuna flagellipes* (Ox-eyed Bean) Seed Flour; An Underutilized Legume in Nigeria. *Am. J. Food Nutr.* 9(1), 49–59.
- Nwaoguikpe RN, Braide W and Ujowundu CO 2011 The Effects of Processing on the Proximate and Phytochemical Compositions of *Mucuna pruriens* seeds (Velvet beans). *Pak. J. Nutr.* 10(10): 947–951.
- Pathania R, Chawla P, Khan H, Kaushik R and Khan MA 2020 An Assessment of Potential Nutritive and Medicinal Properties of *Mucuna pruriens*: a Natural Food Legume. *Biotech* 10(6): 1–15.
- Rakesh B and Praveen N 2020 Biotechnological Approaches for the Production of L- DOPA: A Novel and Potent Anti-Parkinson's Drug from *Mucuna pruriens* (L.) DC. In *Res. Trends Med. Plant Sci.* 10, 181–204.
- Rane M, Suryawanshi S, Patil R, Aware C, Jadhav R, Gaikwad S, Singh P, Yadav S, Bapat V, Gurav R and Jadhav J 2019 Exploring the Proximate Composition, Antioxidant, Anti-Parkinson's and Anti-inflammatory Potential of two Neglected and Underutilized *Mucuna* Species From India. *S. Afr. J. Bot.* 124: 304–310.
- Rai SN, Birla H, Zahra W, Singh S Sen and Singh SP 2017 Immunomodulation of Parkinson's Disease using *Mucuna pruriens* (Mp). *J. Chem. Neuroanatomy*, 85: 27–35.
- Rai SN, Zahra W, Singh S Sen, Birla H, Keswani C, Dilnashin H, Rathore AS, Singh R, Singh R K and Singh SP 2019 Anti-inflammatory Activity of Ursolic Acid in MPTP-Induced Parkinsonian Mouse Model. *Neurotoxicity Res.* 36(3). 452–462.
- Renata LT, Alexandre SS, Ana RNC, Alexandre RPS and Jailane de SA 2015 Nutritional Composition, Phytochemicals and Microbiological Puality of the Legume, *Mucuna pruriens*. *Afr. J. Biotechnol.* 14(8): 676–682.
- Reynolds A, Mann J, Cummings J, Winter N, Mete E and Te Morenga L 2019 Carbohydrate Quality and Human Health: a Series of Systematic Reviews and Meta-analyses. *The Lancet* 393(10170): 434–445.
- Sakamoto K, Khemmarath P, Reiko O and Maro AC 2020 Health , Livelihoods, and Food Intake in Coastal Southeast Tanzania. *J. Faculty Int. Studies, Utsunomiya Univ.* 51: 15–34.
- Sathyanarayana N, Mahesh S, Leelambika M, Jaheer M, Chopra R and Rashmi KV 2012 Genetic Diversity of Wild and Cultivated *Mucuna pruriens* (L.) DC. Accessions Analyzed Using Thirty Morpho-agronomical Characters. *Trop. Subtrop. Agroecosystems* 15(2): 249–259.
- Shan Z, Rehm CD, Rogers G, Ruan M, Wang DD, Hu FB, Mozaffarian D, Zhang FF and Bhupathiraju SN 2019 Trends in Dietary Carbohydrate, Protein, and Fat Intake and Diet Quality among US Adults. *J. Am. Med. Ass.* 322(12): 1178–1187.
- Shimelis EA and Rakshit SK 2005 Proximate Composition and Physico-chemical Properties of Improved Dry Bean (*Phaseolus vulgaris* L.) Varieties Grown in Ethiopia. *Food Sci. Technol.* 38(4): 331–338.
- Siddhuraju P and Becker K 2001 Effect of Various Domestic Processing Methods on Antinutrients and in Vitro Protein and Starch Digestibility of two Indigenous Varieties of Indian Tribal Pulse, *Mucuna pruriens* var. *utilis*. *J. Agric. Food Chem.*,

- 49(6), 3058–3067.
- Siddhuraju P and Becker K 2005 Nutritional and Antinutritional Composition, in Vitro Amino Acid Availability, Starch Digestibility and Predicted Glycemic Index of Differentially Processed *Mucuna* Beans (*Mucuna pruriens* var. *utilis*): an Under-utilised Legume. *Food chem.* 91(2): 275-286.
- Siddhuraju P, Becker K and Makkar HPS 2000 Studies on the Nutritional Composition and Antinutritional Factors of three Different Germplasm Seed Materials of an Under-utilized Tropical Legume, *Mucuna pruriens* var. *Utilis*. *J. Agric. Food Chem.* 48(12): 6048–6060.
- Siddhuraju P, Vijayakumari K and Janardhanan K 1996 Chemical Composition and Protein Quality of the Little-Known Legume, Velvet Bean (*Mucuna pruriens* (L.) DC.). *J. Agric. Food Chem.* 44(9): 2636–2641.
- Siddiq M, Ravi R, Harte JB and Dolan KD 2010 Physical and Functional Characteristics of Selected Dry Bean (*Phaseolus vulgaris* L.) Flours. *Food Sci. Technol.* 43(2): 232–237.
- Suryawanshi SS, Kamble PP, Bapat VA and Jadhav J P 2020 Bioactive Components of Magical Velvet Beans. *Intechopen*. 1–12.
- Timsina J 2018 Can Organic Sources of Nutrients Increase Crop Yields to Meet Global Food Demand? *Agronomy* 214(8): 1–20.
- Tomar S, Chauhan G, Das A and Verma MR 2018 Comparative Evaluation on Phenolic Content and Antioxidant Activity of Legume Sprouts as Affected by Various Solvents for Application in Livestock Products. *Int. J. Curr. Microbiol. Appl. Sci.*, 7(05): 3388–3398.
- Tomé D, Cordella C, Dib O and Péron C 2022 Nitrogen and Protein Content Measurement and Nitrogen to Protein Conversion Factors for Dairy and Soy Protein-based Foods: a Systematic Review and Modelling Analysis. *FAO United Nations* 403–403.
- Tresina PS and Mohan VR 2013 Assessment of Nutritional and Antinutritional Potential of Underutilized Legumes of the Genus *Mucuna*. *Trop. Subtrop. Agroecosystems* 16(2): 155–169. USA National Research Council 1989 Recommended dietary allowances.
- Vadivel V and Janardhanan K 2000 Nutritional and Anti-nutritional Composition of Velvet Bean: An Under-utilized Food Legume in South India. *Int. J. Food Sci. and Nutr.* 51(4): 279–287.
- Wang S, Errington S and Yap HH 2008 Studies on Carotenoids From Lupin Seeds. *Proceedings of the 12th International Lupin Conference, February*, 198–202.
- Wilmot-Dear CM 1984 A Revision of *Mucuna* (Leguminosae-Phaseoleae) in China and Japan. *Kew Bull.* 39(1), 23.