
TANZANIA JOURNAL OF SCIENCE VOLUME 35, 2009

SEROTONIN IMMUNOREACTIVE CELLS AND NERVE FIBERS IN THE MUCOSA OF THE RAT STOMACH

AP Gesase

Department of Anatomy/Histology, Muhimbili University College of Health Sciences
P.O. BOX 65406 Dar es Salaam, Tanzania, Phone 255-22-2150302, Fax: 255-22-2150465
E-mail: agesase@muchs.ac.tz.

ABSTRACT

The current study has showed the morphological features of serotonin (5-hydroxytryptamine) immunoreactivity in the pyloric mucosa of the rat stomach. The immunoreactive elements included the endocrine cells, mast cells and mucosal nerve fibers in the lamina propria. The immunopositive endocrine cells were oval in shape and located mostly in the basal part of the gastric glands. The cell contained a central non-staining area and the peripheral staining zone. The mast cells stained homogenously and appeared to be oval-round in shape. They were located in the upper, middle and basal parts of the mucosa. Serotonin immunoreactive nerve fibers with varicosities were located in the space between the gastric glands in the lamina propria. They appeared not to be in contact with the immunopositive endocrine and mast cells. The current study shows that serotonin may be released by the immunoreactive elements in the stomach and that future work is needed to characterize the ultrastructural features of serotonin positive nerve fibers in the pyloric mucosa.

Keywords: Stomach, Pylorus, Serotonin, Immunohistochemistry, Wistar rats

INTRODUCTION

The gut contains the enteroendocrine cells that stain positive for a number of substances such as gastrin, serotonin (5-hydroxytryptamine), α -Aminobutyric acid (GABA), secretin, substance P (motilin) and cholecystokinin (CCK) that regulate the various activities related to digestion, absorption of food and differentiation and maintenance of the gut epithelium (Johnson 1976; Davanger 1989, Koh *et al.* 1997, Oomori *et al.* 1997, Taniyama *et al.* 2000, Takeda *et al.* 2000, Vincze *et al.* 2001). Gastrin, serotonin and GABA containing cells have been localised in the small and large intestines of various animals and are more numerous in the pyloric antrum of the stomach (Davanger *et al.* 1989; Gesase 2001). GABA- and serotonin-immunopositive have been described to have a similar morphology (Oomori *et al.* 1992).

The cells appear oval or round in shape and many of them have cell processes that extend to reach the lumen of the gastric glands (Oomori *et al.* 1986, Davanger *et al.* 1994, Toole *et al.* 1998). There is limited data on the morphological characterization of serotonin immunopositive cells. One study indicated the presence of nerve fibers and deeply staining and faintly staining cells against serotonin (Yu *et al.* 2001). Such findings need to be clarified and will add knowledge to the morphology of serotonin immunoreactive elements in the stomach.

The current study used the antibody against serotonin to investigate the morphology of serotonin immunoreactivity in the pyloric mucosa of the rat stomach.

MATERIALS AND METHODS

The study was carried out with seven male Wistar rats (about 200g body weight), which

had lived under normal laboratory conditions. The animals received commercial food and water and were kept under constant conditions in the animal house at Muhimbili University College of Health Sciences.

The animals were anaesthetized with ether, thoracotomized, and then perfused via the left cardiac ventricle with 250 ml of physiological saline followed by 150 ml of phosphate-buffered saline (PBS; 0.1 M, pH 7.4) containing 4% paraformaldehyde at 4°C for 10 minutes. The pyloric part of the stomach was removed and cut into small pieces and stored in the same fixative at a temperature of 4°C for 12-24 hours. After rinsing with PBS, the specimens were left overnight in PBS containing 30% sucrose at 4°C for 1 day. The tissues were frozen and cut about 12µm thick using a cryostat, and then mounted on gelatine coated glass slides.

The sections were incubated with a rabbit anti-serotonin (UCB-bioproducts, Brussels, Belgium) diluted 1:10,000 for 24 hours at room temperature. After incubation with primary antibodies the slides were treated with goat biotinylated anti-rabbit IgG and ABC complex (Vector Laboratories, Burlingame, Calif., USA) for 1 hour at room temperature. The antigen-antibody reaction sites were made visible by incubating the sections with diaminobenzidine tetrahydrochloride (DAB) and 0.01% hydrogen peroxide in 2.5mM Tris-HCl buffer (pH 7.6) for 10 minutes. The specificity of the immunohistochemical staining was confirmed by omission of the primary or secondary antibody. No

immunostaining was observed under these conditions through out the experiment.

RESULTS

Observation of the sections revealed serotonin immunoreactive cells and nerve fibers in the mucosa layer of the pylorus (Figs. 1, 2 and 3). The immunoreacted cells were located in the apical, middle and basal parts of the mucosa and close observations revealed the presence of two cell population (Figs. 1 and 2). The first group of cells appeared to have a non-staining central area and the deeply staining peripheral part (Figs. 2 and 3). These cells appeared to be the endocrine cells of the stomach and were located mainly at the basal portion of the mucosa. The endocrine cells were oval in shape, with elongated ends that pointed at the lumina of the gastric glands. The second group of cells that also stained positive against serotonin antibody were randomly distributed in the mucosa of the pylorus, but many of them appeared to be in the middle and upper mucosa (Figs. 2 and 3). These cells stained deeply and uniformly for serotonin antibody and they appeared to be the mast cells in the lamina propria of the stomach wall. They appeared oval-round in shape and did not show cytoplasmic processes. Close observation of the sections stained with serotonin antibody also revealed the presence of immunoreactive nerve fibers in the pyloric mucosa (Fig. 3). Serotonin immunopositive fibers with varicosities were seen to run in the space between the gastric glands and did not appear to establish contact with the immunopositive endocrine cells.




Figure 1: Light micrograph of serotonin (5-hydroxytryptamine) immunoreactivity in the rat stomach showing the mucosa wall (*m*) and immunoreactive cells (*small arrowheads*) that are distributed in the upper, middle and basal parts of the mucosa (*m*). x120



Figure 2: The light micrograph of the rat pyloric mucosa showing the homogenous deeply staining mast cells (*arrowheads*) and endocrine cells (*short thick arrows*) with non-staining central area and the peripheral staining zone. x180




Figure 3: The light micrograph showing the immunopositive nerve fibers (*thin long arrows*) in the rat pyloric mucosa; it also shows the mast cells (*arrowheads*) and immunoreactive endocrine cells (*short thick arrows*). x260

DISCUSSION

The current study has showed the presence of two cell population that stained positive for serotonin antibody. One cell population stained homogenously and appeared to be the mast cells; the endocrine cells showed the central non-staining area and the peripheral staining zone. The results have also showed the presence of serotonin-immunopositive nerve fibers in the pyloric mucosa. The serotonin immunopositive elements may be the source of serotonin that works to regulate various activities in the gut such as smooth muscle contraction and secretion of gastric acids. The mucosal serotonin positive nerve fibers may also play the role of regulating the functions of the epithelia cells in the stomach.

The current results revealed the serotonin immunopositive cells with a central non-staining area that appeared to be the endocrine cells and the deeply homogenously staining mast cells in the

pyloric mucosa. These findings are in agreement with the previous observation in the rat stomach (Yu *et al.* 2001). Other studies also observed the serotonin immunopositive cells but did not report on the presence of deeply staining mast cells (Yamada *et al.* 1999, Bordi *et al.* 2000). These observations indicate that serotonin that is secreted in the gastric wall may come from the endocrine cells and the mast cells; the other source is the immunopositive nerve terminals which have been observed in this study. Serotonin is an important hormone that has been shown to control smooth muscle contraction in the gut and in the stomach it also regulates secretion of gastric acid (Canfield and Spencer 1983, Koop and Arnold 1984, Stephens *et al.* 1989, Wilmer *et al.* 1993, Briejer *et al.* 1997, Takeda *et al.* 2000, Taniyama *et al.* 2000). It has also been shown to play a paracrine role in the stimulation of pancreatic secretion via a vagal cholinergic pathway and may also enhance the growth of

neuritis by stimulating 5-HT_{2B} receptors (Fiorica-Howells *et al.* 2000, Li *et al.* 2001; 2000). Previous work have suggested that the largest source of serotonin to be the endocrine cells of the gut and that mast cells and immunopositive fibers may not have greater contribution to the total amount of serotonin released in the stomach (Schworer *et al.* 1987, Li *et al.* 2001). The current results indicated that both cells showed strong signals for serotonin an indication that they may have an equal role during serotonin secretion. Mast cells are thought to release serotonin during immune response (Askenase *et al.* 1980). It is open for future work if the released serotonin can have other extra immune role.

The results have showed the presence of serotonin immunopositive nerve fibers in the pyloric mucosa. This finding is in agreement with past observations that were made in the gastric mucosa of the rat, guinea pig and golden hamster (Mawe *et al.* 1989, Fujita *et al.* 1995, Fujimiya *et al.* 1997, Toole *et al.* 1998, Chen *et al.* 2001, Yu *et al.* 2001). It has been demonstrated that the cell bodies for the serotonin immunopositive nerve fibers are located in the myenteric plexus (Toole *et al.* 1998, Mawe *et al.* 1989). The role played by the serotonin immunopositive fibers in the stomach is not clearly known. Some authors have suggested that the nerve fibers may exert an important role in the epithelial function of the gastrointestinal tract (Fujimiya *et al.* 1997). Such a suggestion needs to be clarified and is open for future work. Past studies that have been done in other tissues have showed the presence of intraepithelial nerve fibers in the vallate papillae, ejaculatory duct, urethra, tracheobronchial and nasal mucosae that stained positively for calcitonin gene-related peptide (CGRP) immunoreactive and substance P (Lundberg *et al.* 1984, Uddman *et al.* 1985, Yokoyama 1989, Fujita *et al.* 1995, Lee *et al.* 1995, Huang and Lu 1996). It has been suggested that intraepithelial nerve fibers may have a protective role

against noxious stimuli that come from the external environment (Fujita *et al.* 1995). Observations have showed that the gut does not have the intraepithelial nerve fibers that establish contact with the luminal materials (Fujita *et al.* 1995, Chen *et al.* 2001); instead, it contains a large population of the endocrine cells (Yamada *et al.* 1999, Agungpriyono *et al.* 2000, Bordi *et al.* 2000, Vincze *et al.* 2001) that may act as sensor paraneurons in the epithelium that detect noxious materials in the lumen and send the information to the underlying nerve fibers. The serotonin immunopositive nerve fibers observed in the lamina propria may have a role that is yet to be identified. Future work is needed to characterize the ultrastructural features of the serotonin immunopositive nerve fibers in the pyloric mucosa.

ACKNOWLEDGMENTS

I would like to thank staff of the department of Cell biology and Functional Morphology of the Iwate Medical University, Japan for providing me with the antibodies.

REFERENCES

- Agungpriyono S, Macdonald AA, Leus KY, Kitamura N, Adnyane IK, Goodall GP Hondo E and Yamada J 2000 immunohistochemical study on the distribution of endocrine cells in the gastrointestinal tract of the babirusa, *Babirusa babirusa* (Suidae). *Anat. Histol. Embryol.* **29(3)**: 173-178.
- Askenase PW, Bursztajn S, Gershon MD and Gershon RK 1980 T-cell dependent mast cell degranulation and release of serotonin in murine delayed-type hypersensitivity. *J. Exp. Med.* **152**: 1358-1374.
- Bordi C, D'Adda T, Azzoni C and Ferraro G 2000 Classification of gastric endocrine cells at the light and electron microscopical levels. *Microsc. Res. Tech.* **48(5)**: 258-271.
- Briejer MR, Mathis C and Schuurkes JA 1997 5-HT receptor types in the rat ileum longitudinal muscle: focus on 5-

- NT2 receptors mediating contraction. *Neurogastroenterol Motil* **9**(4): 231-237
- Canfield SP and Spencer JE 1983 The inhibitory effects of 5-hydroxytryptamine on gastric acid secretion by the rat isolated stomach. *Br. J. Pharmacol.* **78**: 123-129.
- Chen JJ, Li Z, Pan H, Murphy DL, Tamir H, Koepsell H and Gershon MD 2001 Maintenance of serotonin in the intestinal mucosa and ganglia of mice that lack the high-affinity serotonin transporter: Abnormal intestinal motility and the expression of cation transporters. *J. Neurosci.* **21**(16): 6348-6361.
- Davanger S, Hjelle OP, Babaie E, Larsson L-I, Hougaard D, Storm-Mathisen J and Ottersen OP 1994 Colocalization of γ -aminobutyrate and gastrin in the rat antrum: An immunocytochemical and in situ hybridization study. *Gastroenterol.* **107**: 137-148.
- Davanger S, Ottersen OP, Storm-Mathisen J 1989 GABA-immunoreactive cells in the rat gastrointestinal epithelium. *Anat. Embryol.* **179**(3): 221-116.
- Fiorica-Howells E, Maroteaux L and Gershon MD 2000 serotonin and the 5-HT_{2B} receptor in the development of enteric neurons. *J. Neurosci.* **20**: 294-305.
- Fujimiya M, Okumiya K, Yamane T and Maeda T 1997 Distribution of serotonin-immunoreactive nerve cells and fibers in the rat gastrointestinal tract. *Histochem. Cell. Biol.* **107**(2): 105-114.
- Fujita T, Iwanaga T, Lee S-H and Terada M 1995 Sensor cells and nerve terminals in some gut-derived organs: a review. *IT J. Anat Embryol.* **100**(1): 197-204.
- Gesase AP 2001 Gamma aminobutyric acid immunoreactivity in the rat stomach during postnatal development. *T. Med. J.* **16**(3): 31-34.
- Huang YJ and Lu KS 1996 Immunohistochemical studies on protein gene products 9.5, serotonin and neuropeptides in vallate taste buds and related nerves of the guinea pig. *Arch. Histol. Cytol.* **59**(5): 433-441.
- Johson LR 1976 The trophic action of gastrointestinal hormones. *Gastroenterology* **70**: 278-288.
- Koh TJ, Goldenring JR, Ito S, Mashimo H, Kopin AS, Varro A, Dockray GJ and Wang TC 1997 Gastrin deficiency results in altered gastric differentiation and decreased colonic proliferation in mice. *Gastroenterology* **113**(3): 1015-1025.
- Koop H and Anold R 1984 Serotonergic control of somatostatin and gastrin release from the isolated rat stomach. *Regul Pept* **9**: 101-108
- Lee SH, Iwanaga T and Fujita T 1995 Intraepithelial nerve fibers in the nasal mucosa of the rat with special reference to the localization of CGRP, VIP and nitric oxide (NO). *Arch. Histol. Cytol.* **58**(4): 449-456
- Li Y, Hao Y and Owyang C 2000 Serotonin released from intestinal enterochromaffin cells mediates luminal and non-cholecystokinin-stimulated pancreatic secretion in rats. *Gastroenterology* **118**: 1197-1207.
- Li Y, Wu XY, Zhu JX and Owyang C 2001 Intestinal serotonin acts as paracrine substance to mediate pancreatic secretion stimulated by luminal factors. *Am. J. Physiol. Gastrointest. Liver Physiol.* **281**: G916-G923.
- Lundberg JM, Hokfelt T, Martling CR, Saria A and Cuello C 1984 Substance P-immunoreactive sensory nerves in the lower respiratory tract of various mammals including man. *Cell Tissue Res.* **235**: 251-261.
- Mawe GM, Schemann M, Wood JD and Gershon MD 1989 Immunocytochemical analysis of potential neurotransmitters present in the myenteric plexus and muscular layers of the corpus of the guinea pig stomach. *Anat. Rec.* **224**(3): 431-442.
- Oomori Y 1986 Immunohistochemistry and morphometry of gastrin cells in the rat pyloric antrum during starvation. *Anat. Embryol.* **175**: 7-14.

- Oomori Y, Iuchi H, Ishikawa K, Satoh Y and Ono K 1992 Gamma-aminobutyric acid immunoreactivity in the enterochromaffin cells of the rat stomach. *Acta. Anat.* **145**: 127-131.
- Oomori Y, Satoh Y, Ishikawa K and Gesase AP 1997 Exocytosis in the antral gastrin cells of mouse, rat and guinea pig after stimulation by carbamylcholine. *Cell Tissue Res.* **289**: 463-472.
- Schwörer H, Racké K and Kilbinger H 1987 Spontaneous release of endogenous 5-hydroxytryptamine and 5-hydroxyindoleacetic acid from the isolated vascularly perfused ileum of the guinea pig. *Neuroscience* **21**: 297-303.
- Stephens RL, Garrick T, Weiner H and Tache Y 1989 serotonin depletion potentiates gastric secretory and motor responses to vagal but not peripheral gastric stimulants. *J. Pharmacol. Exp. Ther.* **251**: 524-530.
- Takeda M, Tsukamoto K, Sakurai-Yamashita Y, Suzuki T and Taniyama K 2000 Facilitation of acetylcholine release by SK-951, a benzofuran derivative, via the 5-hydroxytryptamine₄ receptor in guinea pig stomach. *Jpn. J. Pharmacol.* **82(2)**: 138-143.
- Taniyama K, Makimoto N, Furuichi A, Sakurai-Yamashita Y, Nagase Y, Kaibara M and Kanematsu T 2000 Functions of peripheral 5-hydroxytryptamine receptors, especially 5-hydroxytryptamine₄ receptor, in gastrointestinal motility. *J. Gastroenterol.* **35(8)**: 575-582.
- Toole L, Belai A and Burnstock G 1998 A neurochemical characterisation of the golden hamster myenteric plexus. *Cell Tissue Res.* **291(3)**: 385-394.
- Uddman R, Luts A and Sundler F 1985 Occurrence and distribution of calcitonin gene-related peptide in the mammalian respiratory tract and middle ear. *Cell Tissue Res.* **241**: 551-555.
- Vincze E, Kantor O, Kausz M, Nemeth J, Arimura A, Gonda P and Kovacs K 2001 Comparative study on the appearance of various bioactive peptides in foregut derivatives during the ontogenesis. *J. Physiol. Paris* **95(1-6)**: 99-103.
- Wilmer A, Tack J, Coremans G, Janssens J, Peeters T and Vantrappen G 1993 5-hydroxytryptamine-3 receptors are involved in the initiation of gastric phase-3 motor activity in humans. *Gastroenterology* **105**: 773-780.
- Yamada J, Tauchi M, Rerkamnuaychoke W, Endo H, Chungsamarnyart N, Kimura J, Kurohmaru M, Hondo E, Kitamura N, Nishida T and Hayashi Y 1999 Immunohistochemical survey of the gut endocrine cells in the common tree shrew (*tupaia belangeri*). *J. Vet. Med. Sci.* **61(7)**: 761-767.
- Yokoyama R 1989 Histochemical and immunohistochemical studies on intraepithelial nerve fibers in the ejaculatory duct of the monkey (*Macaca fuscata*). *Kurume Med. J.* **36(3)**: 101-112.
- Yu PL, Fujimura M, Hayashi N, Nakamura T, and Fujimiya M 2001 Mechanisms in regulating the release of serotonin from the perfused rat stomach. *Am. J. Physiol. Gastrointest. Liver Physiol.* **280**: 1099-1105.

